




The ecosulfur Advantage

- ✓ **Minimal Dust:** A very uniform, free-flowing water dispersible granule vs. standard wettable powder formulations
- ✓ **Ecosulfur is compatible with:**
 - Most fungicides – organic and copper compounds, including Bordeaux mixture
 - Most insecticides, herbicides and miticides
 - Emulsifiable concentrates and spray oils
- ✓ **Excellent Particle size** range from 2 to 5 microns (median 3 microns) for better coverage and increased effectiveness against fungal diseases
- ✓ **Excellent Suspensibility:** In both hard and soft water the suspensibility is the same with 80% remaining in suspension after one hour. Minimal agitation needed to re-suspend
- ✓ **Wettability:** Instantaneous when added to water
- ✓ **Input for Organic Production:** Registered for use in organic farming systems. ACO & NOP Approved
- ✓ **Instantaneous Dispersion:** Dissolves completely in water, within seconds of adding to tank with little to no residue left in the spray tank (see images below)



ecosulfurTM

800 WG ORGANIC* FUNGICIDE & MITICIDE

ACTIVE CONSTITUENT: 800G/KG SULFUR (S) PRESENT AS ELEMENTAL SULFUR

APPLICATION RATES

TREE AND VINE CROPS

CROP	DISEASE	STATE	RATE/100L	CRITICAL COMMENTS
Citrus	Citrus Rust Mite, Brown Citrus Rust Mite, Bud Mite, Moss, Lichens	NSW, Vic, Qld, SA, WA only	400-500g	Apply during July-August. Leave at least 3 weeks between sprays of oil and product.
	Melanose	NSW, WA only	400-500g	Will assist control when applied late August.
	White Louse Scale	NSW, WA only	200-300g	May be added to copper spray at petal fall or applied separately during Spring or Autumn.
Grapes Table Grapes	Vine Mite, Grapeleaf Blister Mite	NSW, Vic, SA, WA, TAS only	500g	Apply before sprouting. Ensure thorough coverage.
	Vine Mite	Qld only	500g	Apply before sprouting. Ensure thorough coverage.
	Powdery Mildew (<i>Oidium</i> spp.) Mites	NSW, Vic, SA, WA, TAS only	100-200g	Apply immediately after budburst, then every 2-3 weeks or as required.
Grapevines Wine Grapes only	Vine Mite, Grapeleaf Blister Mite	NSW, Vic, SA, WA, TAS only	500g	Apply at late budburst. During summer, build up of mites is checked by sulfur sprays for powdery mildew. An application immediately after picking may aid control in vineyards where mites are a severe problem.
	Vine Mite	Qld only		
	Powdery Mildew (<i>Oidium</i> spp.) Mites	NSW, Vic, SA, WA, TAS only	200-600g	Apply when shoots are 20 to 25cm long and repeat at intervals of 10 to 14 days during humid weather. Use the higher rates when disease pressure is high and/or a higher level of control is required. Crop phytotoxicity may occur when applications are made during hot conditions. DO NOT apply during the heat of the day. Use the higher rates early in the season before the temperatures rise. DO NOT apply more than 6kg per hectare.
Kiwifruit	Twospotted Mite	NSW, WA only	150-200g	Spray only when the pest is present. Repeat application only when necessary.
Mangosteens	Powdery Mildew Mites	All States	400g	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.
Peaches, Nectarines, Plums	Rust	NSW, Vic, SA, Tas only	200g	Apply as required during November to January.
		Qld only	350g	Apply at petal fall only.
	Brown Rot (<i>Monilinia fructicola</i>) Blossom Blight (<i>Monilinia laxa</i>)	NSW, Vic, SA, WA, TAS only	200g	Apply at petal fall and then as a cover spray at regular intervals beginning 3-4 weeks after petal fall.
		Qld only	350g	Apply at petal fall only.
Pome Fruit	Powdery Mildew	All States	400g	Start spraying early. Apply at least 2 sprays before blossom. Apply every 2 weeks after blossom or as required. Regular sprays will also aid in the control of Black Spot (Scab).
		Vic, SA, TAS only	200g	
	Powdery Mildew, Black Spot (Scab)	Qld only	350g	
Rambutans	Powdery Mildew, Mites	All States	400g	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.

OTHER CROPS

Ornamentals	Powdery Mildew Twospotted Mite	Vic, SA, WA, TAS only	200g	Apply when pest or disease is first seen and repeat every 10-21 days.
		Qld only	350g	
	Powdery Mildew	NSW only	200g	
Strawberries	Powdery Mildew	NSW only	200-300g	Apply when the disease is first noticed then at 10-14 day intervals during humid weather.
Tomatoes for processing	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite, Twospotted Mite	Vic, SA, WA, TAS only	2 kg/ha	Apply when pest or disease is first seen and repeat as necessary. Crop phytotoxicity may occur when applications occur during hot conditions. DO NOT apply during the heat of the day.
	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite	NSW only		
Vegetables (except Cucumbers or Rockmelons)	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite, Twospotted Mite	Vic, SA, WA, TAS only	200g	
	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite	NSW only	200g	
	Powdery Mildew, Bean Rust, Tomato Russet Mite	Qld only	300g	



ORGANIC CROP PROTECTANTS
 1956 Dandenong Rd, Clayton VIC 3168
 Telephone: 1800 634 204 (Free Call)
 ocp.com.au

